Discover and Verify

The sum of the powers of the first n positive integers gives a list of interesting series:

$$1 + 2 + \dots + n = \frac{n^2 + n}{2} = \frac{n(n+1)}{2} \dots (1)$$

$$1^2 + 2^2 + \dots + n^2 = \frac{n^3}{3} + \frac{n^2}{2} + \frac{n}{6} = \frac{n(n+1)(2n+1)}{6} \dots (2)$$

$$1^3 + 2^3 + \dots + n^3 = \frac{n^4}{4} + \frac{n^3}{2} + \frac{n^2}{4} = \left[\frac{n(n+1)}{2}\right]^2 \dots (3)$$

$$1^4 + 2^4 + \dots + n^4 = \frac{n^5}{5} + \frac{n^4}{2} + \frac{n^3}{3} - \frac{n}{30} = \frac{n(n+1)(2n+1)(3n^2 + 3n - 1)}{30} \dots (4)$$

The pursuit of the coefficients in the formulas ended up in the rather old story called the Faulhaber's theorem. The Johann Faulhaber's formula expresses the sum of the k-th powers of the first n positive integers.

$$\sum_{i=0}^{n} i^{k} = 1^{k} + 2^{k} + \dots + n^{k}$$

$$\sum_{i=1}^{n} i^{k} = \frac{1}{k+1} \sum_{i=1}^{k} (-1)^{j} {k+1 \choose j} B_{j} n^{k+1-j}$$

where $B_1 = -\frac{1}{2}$ and B_j 's are Bernoulli numbers

The understanding and proof are remote for most secondary students and you may forget about it.

However, here is a mysterious pattern I came across for finding the coefficients you may enjoy.

First, study carefully the Coefficient Triangle below:

					1 ₁						
				1 ₁		12					
			11		3 ₂		2 ₃				
		11		7 ₂		12 ₃		64			
	11		15 ₂		50 ₃		60 ₄		24 ₅		
1 ₁		31 ₂		130 ₃		390 ₄		365 ₅		120 ₆	

Task 1 Discover how the numbers in the Coefficient Triangle are constructed.

Hints: 1. Study the subscripts in red. How to put in the numbers?

2. How to find the numbers in black? Recall the Pascal triangle in Binomial Theorem, but here you need to do some multiplications.

If we write : $\sum_{i=0}^n i^k = 1^k + 2^k + \dots + n^k$ and C_r^n be the Binomial coefficients, we get:

$$\sum_{i=1}^{n} i^0 = 1C_1^n \quad ... (5)$$

$$\sum_{i=1}^{n} i^{1} = 1C_{1}^{n} + 1C_{2}^{n} \dots (6)$$

$$\sum_{i=1}^{n} i^2 = 1C_1^n + 3C_2^n + 2C_3^n \dots (7)$$

$$\sum_{i=1}^{n} i^{3} = 1C_{1}^{n} + 7C_{2}^{n} + 12C_{3}^{n} + 6C_{4}^{n} \dots (8)$$

$$\sum_{i=0}^{n} i^4 = 1C_1^n + 15C_2^n + 50C_3^n + 60C_4^n + 24C_5^n \dots (9)$$

$$\sum_{i=0}^{n} i^{5} = 1C_{1}^{n} + 31C_{2}^{n} + 130C_{3}^{n} + 390C_{4}^{n} + 365C_{5}^{n} + 120C_{6}^{n} \dots (10)$$

Note that the coefficients in the formulas (5) - (10) are just the numbers (in black) in the Coefficient Triangle above.

Task 2 Checking

Show that the equations are correct.

$$(7) = (2), \qquad (8) = (3), \qquad (9) = (4)$$

The general proof that this pattern works well for equation (10) or beyond may not be easy and is not discuss here. However, readers may try the followings.

Task 3 Verification

- 1. Prove that equation (7) or (2) is correct by Mathematical Induction.
- **2.** Prove that equation (8) or (3) is correct by considering the identity:

$$i^{2}(i+1)^{2} - (i-1)^{2}i^{2} = 4i^{3}$$

3. Prove that equation (9) or (4) is correct by considering the identity:

$$(i+1)^5 - i^5 = 5i^4 + 10i^3 + 10i^2 + 5i + 1$$